



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTION OF A PROBLEM.

BY ISAAC H. TURRELL, CUMMINSVILLE, OHIO.

Given three circles touching each other externally; if another group of three be drawn touching each other in like manner, and so situated that each circle of either group touches two of the other, to prove that the square of the common tangent of any two of the circles, which do not touch each other, is equal to twice the product of their diameters.

Solution. This property is easily proved by means of the following principle, due to Mr. Casey, and found on page 113, Salmon's Conics, Fourth Edition: "If four circles be all touched by the same fifth circle, the lengths of their common tangents are connected by the following relation:

$\overline{12} \cdot \overline{34} \pm \overline{14} \cdot \overline{23} \pm \overline{13} \cdot \overline{24} = 0$, where $\overline{12}$ denotes the length of a common tangent to the first and second circles, &c."

Denoting the circles of the first group and their radii by a, b, c , and of the second by x, y, z , there will be three pairs of non-touching circles, viz; a, x , b, y , and c, z . Let T, T', T'' be their common tangents, then as z is touched by a, b, x, y , which touch each other consecutively,

$$2\sqrt{(ab)} \times 2\sqrt{(xy)} + 2\sqrt{(ay)} \times 2\sqrt{(bx)} = T.T',$$

or $8\sqrt{(abxy)} = T.T'.$

Similarly $8\sqrt{(bcyz)} = T'.T'',$

and $8\sqrt{(cazx)} = T''.T,$ whence, by elimination,

$$T^2 = 8ax = \text{twice the product of the diameters of } a \text{ and } x.$$

NOTE.—TO THE STUDENT OF PHYSICAL ASTRONOMY: When you undertake the study of the theory of Perturbations of the planetary motions, do not take up any extended work on the subject and attempt to master all its details at one reading, but try to get the *general principles* of the subject — not the most general methods—and as soon as you have done that, commence to prepare with your own hands a treatise on the subject for your own use. Continue to *think* on the subject till you can explain to another all the details so far as you go, and do this in your treatise. By all means commence with the *Lunar Theory*, and I should say the "*Variation of Constants*".

DAVID TROWBRIDGE.